





Рис. 2. Продукты ПЦР на матрице кДНК:

1–3, 11, 12 – фрагменты (~1000 п. н.), соответствующие *acdS*-гену (1 – Cu^{2+} ; 2 – Cr^{6+} ; 3 – Pb^{2+} ; 11, 12 – 200 ммоль/л NaCl); 4, 7, 13, 20 – 1 kb DNA Ladder; 5, 19 – pBI121/*acdS*-B37; 6, 14, 15 – негативный контроль; 8–10, 17, 18 – фрагменты (~500 п. н.), соответствующие гену *Ef-1a* (8 – Cu^{2+} ; 9 – Cr^{6+} ; 10 – Pb^{2+} ; 17, 18 – 200 ммоль/л NaCl); 16 – *GAPDH*

Fig. 2. PCR products on the cDNA matrix:

1–3, 11, 12 – fragments (~1000 bp) corresponding to the *acdS*-gene (1 – Cu^{2+} ; 2 – Cr^{6+} ; 3 – Pb^{2+} ; 11, 12 – 200 mmol/L NaCl); 4, 7, 13, 20 – 1 kb DNA Ladder; 5, 19 – pBI121/*acdS*-B37; 6, 14, 15 – negative control; 8–10, 17, 18 – fragments (~500 bp) corresponding to the gene *Ef-1a* (8 – Cu^{2+} ; 9 – Cr^{6+} ; 10 – Pb^{2+} ; 17, 18 – 200 mmol/L NaCl); 16 – *GAPDH*