



УДК 511.42

ВКЛАД ЙОНАСА КУБИЛЮСА В МЕТРИЧЕСКУЮ ТЕОРИЮ ДИОФАНТОВЫХ ПРИБЛИЖЕНИЙ ЗАВИСИМЫХ ПЕРЕМЕННЫХ

В. В. БЕРЕСНЕВИЧ¹⁾, В. И. БЕРНИК²⁾,
Ф. ГЁТЦЕ³⁾, Е. В. ЗАСИМОВИЧ²⁾, Н. И. КАЛОША²⁾

¹⁾Колледж им. Джеймса Рашолма, Йоркский университет, Западный кампус,
YO10 5DD, г. Йорк, Великобритания

²⁾Институт математики НАН Беларуси, ул. Сурганова, 11, 220072, г. Минск, Беларусь

³⁾Билефельдский университет, Университетситрассе, 25, D-33615, г. Билефельд, Германия

Посвящается 100-летию со дня рождения академика Йонаса Кубилюса, который является основоположником метрической теории диофантовых приближений. Проводится обзор наиболее важных результатов, полученных в метрической теории диофантовых приближений. Отмечается, что за последние 70 лет в области диофантовых приближений сделано много выдающихся достижений. Упоминаются работы лауреатов Филдсовской премии Алана Бейкера и Григория Маргулиса, а также ученика Й. Кубилюса, академика АН БССР Владимира Спринджука, который в 1964 г. решил известную проблему Малера и стал основателем белорусской школы теории чисел.

Ключевые слова: Й. Кубилюс; диофантовы приближения; проблема Малера; метрическая теория чисел; трансцендентные и алгебраические числа.

Образец цитирования:

Бересневич ВВ, Берник ВИ, Гётце Ф, Засимович ЕВ, Калоша НИ. Вклад Йонаса Кубилюса в метрическую теорию диофантовых приближений зависимых переменных. *Журнал Белорусского государственного университета. Математика. Информатика.* 2021;3:34–50 (на англ.).
<https://doi.org/10.33581/2520-6508-2021-3-34-50>

For citation:

Beresnevich VV, Bernik VI, Götze F, Zasimovich EV, Kalosha NI. Contribution of Jonas Kubilius to the metric theory of Diophantine approximation of dependent variables. *Journal of the Belarusian State University. Mathematics and Informatics.* 2021; 3:34–50.
<https://doi.org/10.33581/2520-6508-2021-3-34-50>

Авторы:

Виктор Вячеславович Бересневич – доктор физико-математических наук, профессор; профессор.

Василий Иванович Берник – доктор физико-математических наук, профессор; главный научный сотрудник отдела теории чисел.

Фридрих Гётце – доктор физико-математических наук, профессор; профессор.

Елена Васильевна Засимович – аспирантка отдела теории чисел. Научный руководитель – В. И. Берник.

Николай Иванович Калоша – кандидат физико-математических наук; старший научный сотрудник отдела теории чисел.

Authors:

Victor V. Beresnevich, doctor of science (physics and mathematics), full professor; professor.

victor.beresnevich@york.ac.uk

Vasily I. Bernik, doctor of science (physics and mathematics), full professor; chief researcher at the department of number theory.

bernik.vasili@mail.ru

Friedrich Götze, doctor of science (physics and mathematics), full professor; professor.

goetze@math.uni-bielefeld.de

Elena V. Zasimovich, postgraduate student at the department of number theory.

elena.guseva.96@yandex.by

Nikolai I. Kalosha, PhD (physics and mathematics); senior researcher at the department of number theory.

kalosha@im.bas-net.by





CONTRIBUTION OF JONAS KUBILIUS TO THE METRIC THEORY OF DIOPHANTINE APPROXIMATION OF DEPENDENT VARIABLES

V. V. BERESNEVICH^a, V. I. BERNIK^b,
F. GÖTZE^c, E. V. ZASIMOVICH^b, N. I. KALOSHA^b

^aJames College, University of York, Campus West,
YO10 5DD, York, United Kingdom

^bInstitute of Mathematics, National Academy of Sciences of Belarus,
11 Surhanava Street, Minsk 220072, Belarus

^cBielefeld University, 25 Universitätsstraße, Bielefeld D-33615, Germany

Corresponding author: N. I. Kalosha (kalosha@im.bas-net.by)

The article is devoted to the latest results in metric theory of Diophantine approximation. One of the first major result in area of number theory was a theorem by academician Jonas Kubilius. This paper is dedicated to centenary of his birth. Over the last 70 years, the area of Diophantine approximation yielded a number of significant results by great mathematicians, including Fields prize winners Alan Baker and Grigori Margulis. In 1964 academician of the Academy of Sciences of BSSR Vladimir Sprindžuk, who was a pupil of academician J. Kubilius, solved the well-known Mahler's conjecture on the measure of the set of S -numbers under Mahler's classification, thus becoming the founder of the Belarusian academic school of number theory in 1962.

Keywords: J. Kubilius; Diophantine approximation; Mahler's conjecture; metric number theory; transcendence and algebraic numbers.

*Dedicated to the centenary of
academician Jonas Kubilius's birth*

Introduction

Academician Jonas Kubilius devoted his life to research in theory of probability and number theory. He was one of the founders of metric theory of Diophantine approximation, obtaining one of the earliest major results in this field [1; 2] and influencing the work of his pupil Vladimir Sprindžuk, who in 1964 proved the famous Mahler's conjecture [3–5].

First results in metric theory of Diophantine approximation were obtained by Émile Borel in the beginning of the 20th century [6], and they were later significantly improved in a seminal work of Alexander Khintchine [7].

Let $\psi(x)$ be a monotonic decreasing function of $x > 0$, and let μB denote the Lebesgue measure of a measurable set $B \subset \mathbb{R}$. Let $\mathcal{L}_1(\psi)$ denote the set of real numbers in the interval $I \subset \mathbb{R}$ such that the inequality

$$\left| x - \frac{p}{q} \right| < \frac{\psi(q)}{q} \quad (1)$$

has infinitely many solutions in integers $p \in \mathbb{Z}$ and positive integers $q \in \mathbb{N}$.

Theorem 1 (Khintchine's theorem). *The Lebesgue measure of the set $\mathcal{L}_1(\psi)$ satisfies*

$$\mu \mathcal{L}_1(\psi) = 0 \text{ if } \sum_{q=1}^{\infty} \psi(q) < \infty, \quad (2)$$

$$\mu \mathcal{L}_1(\psi) = \mu I \text{ if } \sum_{q=1}^{\infty} \psi(q) = \infty. \quad (3)$$

Note that in the case of convergence (2) the theorem also holds without the monotonicity requirement on the function $\psi(x)$. Khintchine's theorem was later generalised by A. V. Groshev for system of linear forms [8].

One important generalisation of the above setting considered by A. Khintchine concerns small values of integral polynomials, see articles [3; 5] and monographs [4; 9–11]. In particular, in [12] A. Khintchine proved the following theorem.



Theorem 2. For all $\varepsilon > 0$ and an arbitrary interval I , the inequality

$$|P(x)| < \varepsilon H^{-n} \quad (4)$$

has infinitely many solutions in polynomials

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0, P(x) \in \mathbb{Z}[x],$$

of degree n and height $H = H(P) = \max_{0 \leq j \leq n} |a_j|$ for almost all points $x \in I$.

Clearly, we can find $x \in \mathbb{R}$ such that for $\varepsilon = \varepsilon(x)$ the inequality opposite to (4) is satisfied. Taking, for example, $x_1 = \sqrt[n+1]{2}$, we have

$$|P(x_1)| > c(n)H^{-n}$$

for an appropriate positive constant $c(n)$.

The inequality (4) can be viewed as the first result in metric Diophantine approximation of dependent variables since it relates to approximation of zero by values of an integer polynomial of an arbitrary degree.

In what follows, $c_1 = c_1(n)$, c_2, \dots will denote quantities that depend on n and do not depend on H . Quantities A and B satisfy the inequality $A \ll B$, where \ll is the Vinogradov symbol, if there exists a constant c such that $A < cB$.

In 1932 Kurt Mahler [13] proposed his classification of real and complex numbers based on the behaviour of best polynomial approximations of zero $P(\xi)$ and $P(z)$ as the height H tends to infinity at respectively real and complex points $\xi \in \mathbb{R}, z \in \mathbb{C}$. A detailed description of Mahler's classification can be found in Sprindžuk's [3; 4], Schmidt's [14] and Bugeaud's [15] monographs. K. Mahler divided the set of real numbers into four classes, namely the A -, S -, T - and U -numbers. He formulated the famous Mahler's conjecture, which became the main problem in metric theory of Diophantine approximations for several decades.

Let $M_n(w)$ be the set of real numbers $x \in I$ such that the inequality

$$|P(x)| < H^{-w}, w > 0,$$

has infinitely many solutions in polynomials $P(x) \in \mathbb{Z}[x]$, $\deg P = n$.

Conjecture (Mahler's conjecture). For $w > n$ we have $\mu M_n(w) = 0$.

For complex numbers, conjecture 1 can be formulated as follows: define the set $\mathcal{K}_n(v)$ of $z \in \mathbb{C}$ such that the inequality $|P(z)| < H^{-v}$ has infinitely many solutions in $P(z) \in \mathbb{Z}[z]$, then for $v > \frac{n-1}{2}$ we have $\mu_2 \mathcal{K}_n(v) = 0$, where μ_2 is the two-dimensional Lebesgue measure on the complex plane \mathbb{C} .

Mahler's problem can also be formulated in terms of simultaneous Diophantine approximation: let $S_n(t)$, $t > 0$, be the set of real numbers x such that the inequality

$$\max_{1 \leq l \leq n} |qx^l - p_l| < q^{-t}, t > n^{-1}, \quad (5)$$

has infinitely many solutions in integer vectors $\bar{m} = (q, p_1, \dots, p_n)$, then $\mu S_n(t) = 0$.

The two formulations are equivalent by Khintchine's transference principle [10]; the equality $\mu M_n(w) = 0$ implies that $\mu S_n(t) = 0$ and vice versa.

K. Mahler was able to prove that $\mu M_n(w) = 0$ for $w > 4n$, with other researchers offering consecutive improvements: $w > 3n$ by Jurjen Koksma [16], $w \leq 2$ by William LeVeque [17]; $w \leq 2 - \frac{2}{n}$ by Friedrich Kasch and Bodo Volkmann [18]; $w \leq 2 - \frac{7}{3n}$ by Wolfgang Schmidt [19]; $w \leq \frac{3}{2}$, $w \leq \frac{4}{3}$ by B. Volkmann [20] using Davenport's lemma [21].

The first proof of Mahler's conjecture for $n = 2$ was obtained by academician J. Kubilius for the inequality (5) using the method developed by academician Ivan Vinogradov [1; 22].

From Minkowski's linear forms theorem we see that the system of Diophantine inequalities

$$\max(\|xq\|, \|x^2q\|) < q^{-1/2}$$

has infinitely many solutions for all $x \in \mathbb{R}$ and positive integers $q \in \mathbb{N}$. However, if we slightly reduce the order of the right-hand side to obtain

$$\max(\|xq\|, \|x^2q\|) < q^{-1/2-\varepsilon},$$



then this new inequality has, for an arbitrarily small positive ε , only a finite number of solutions for almost all x . This result was proved by J. Kubilius in 1949 [1], thus proving Mahler's conjecture in the quadratic case. Soon thereafter John William Scott Cassels [23] obtained an improvement of Kubilius's result, proving that the system of inequalities

$$\|xq\| < \varphi(q), \|x^2q\| < f(q)$$

has a finite number of solutions in $q \in \mathbb{N}$ for almost all x if the series

$$\sum_{q=1}^{\infty} \varphi(q)f(q)$$

converges and

$$f(q) \geq \max(\varphi(q), q^{-1/2} \log q \sigma(q)),$$

where $\sigma(q)$ is the number of positive integer divisors of q . In 1959 J. Kubilius improved this result by relaxing the requirements on $\max(\varphi(q), f(q))$, proving the following theorem [2].

Theorem 3. *The inequality*

$$\max(\|xq\|, \|x^2q\|) < \psi(q)$$

has only a finite number of solutions in $q \in \mathbb{N}$ for a positive function $\psi(q)$ if $q^{-1/2} \psi(q)$ is non-increasing and the series

$$\sum_{q=1}^{\infty} q^{-1/2} \psi(q)$$

converges.

In a private conversation with the V. Bernik, J. Kubilius mentioned that he had obtained the proof of Leveque's result prior to the publication of Leveque's paper, but J. Kubilius wasn't expedient in publishing this proof.

Mahler's problem was solved in 1964 by Belarusian mathematician V. Sprindžuk, a pupil of J. Kubilius. V. Sprindžuk proved Mahler's conjecture not only in the fields \mathbb{R} and \mathbb{C} , but also reformulated and proved it for p -adic numbers and formal power series. V. Sprindžuk laid down the groundworks of metric theory of Diophantine approximation, publishing two monographs in Russian and English [4; 11].

The next few sections of this article will be devoted to solutions and generalisations of the problems that were posed in the 1950–60s and were related to Koksma's and Mahler's classifications, as well as the classical problems of Vladimir Sprindžuk, Alan Baker and Wolfgang Schmidt [24]. After that, we'll move on to applications of the methods of metric theory of Diophantine approximation to quantifying distributions of rational and algebraic numbers, as well as discriminants and resultants of integer polynomials.

To conclude the article, we will touch upon applications of Diophantine approximation in mathematical physics and wireless communications.

Generalisations of the Mahler – Sprindžuk problem

Several results related to Mahler's conjecture have been improved and generalised. In particular, a full analogue of Khintchine's theorem was proved for the inequalities (2) and (3). Let $\mathcal{L}_n(\psi)$ denote the set of $x \in I$ such that the inequality

$$|P(x)| < H^{-n+1} \psi(H)$$

has infinitely many solutions in polynomials $P(x) \in \mathbb{Z}[x]$ of degree n and height $H = H(P)$.

Theorem 4. *The Lebesgue measure of the set $\mathcal{L}_n(\psi)$ is*

$$\mu \mathcal{L}_n(\psi) = 0 \text{ if } \sum_{H=1}^{\infty} \psi(H) < \infty, \quad (6)$$

$$\mu \mathcal{L}_n(\psi) = \mu I \text{ if } \sum_{H=1}^{\infty} \psi(H) = \infty. \quad (7)$$

The equality (6) was proved by Vasili Bernik in the paper [25], and (7) was proved by Victor Beresnevich in [26]. In the case of convergence (6), theorem 4 also holds if the monotonicity requirement on $\psi(H)$ is omitted [27], as will be discussed later on. Analogues of theorem 4 also hold in the complex case [28], the p -adic



case [29; 30], and in the case of approximation by algebraic numbers [15; 31; 32]. Many of these results have become parts of monographs [4; 11; 33].

Let $f_1(x), \dots, f_n(x) \in C^{n+1}(I)$ be $n + 1$ times continuously differentiable functions of the real variable $x \in I$ such that their Wronskian is non-zero,

$$W(x) = \begin{vmatrix} f_1'(x) & f_2'(x) & \dots & f_n'(x) \\ f_1''(x) & f_2''(x) & \dots & f_n''(x) \\ \dots & \dots & \dots & \dots \\ f_1^{(n)}(x) & f_2^{(n)}(x) & \dots & f_n^{(n)}(x) \end{vmatrix} \neq 0, \quad (8)$$

for almost all $x \in I$. Let $\mathcal{K}_n(\psi)$ denote the set of $x \in I$ such that the inequality

$$|F(x)| = |a_n f_n(x) + \dots + a_1 f_1(x) + a_0| < H^{-n+1} \psi_1(H)$$

has infinitely many solutions, where H is the «height» of the function $F(x)$ defined as $\max_{0 \leq j \leq n} |a_j|$. W. Schmidt [19] proved that for $\psi_1(H) = H^{-\gamma}$, $\gamma > 1$, we have the equality $\mu \mathcal{K}_2(\psi_1) = 0$. The paper [34] proves this result for $n = 3$.

V. Sprindžuk conjectured [4] that for an arbitrary n and $\psi_1(H) = H^{-\gamma}$, $\gamma > 1$, we have $\mu \mathcal{K}_n(\psi_1) = 0$. This conjecture was proved by Dmitry Kleinbock and Grigori Margulis [35]. Soon thereafter, the following theorem was proved for a curve $S(x) = (f_1(x), \dots, f_n(x))$ satisfying the condition (8).

Theorem 5. *The measure of $\mathcal{K}_n(\psi)$ is given as*

$$\mu \mathcal{K}_n(\psi) = 0 \text{ if } \sum_{H=1}^{\infty} \psi(H) < \infty, \quad (9)$$

$$\mu \mathcal{K}_n(\psi) = \mu I \text{ if } \sum_{H=1}^{\infty} \psi(H) = \infty. \quad (10)$$

The equality (9) was independently proved by V. Beresnevich [36] and V. Bernik, D. Kleinbock and G. Margulis [37], and (10) was proved by the latter authors in [38]. Several related results in metric theory of Diophantine approximation are presented in monographs [4; 11; 15; 33; 39–41] and the article [42].

Diophantine approximation and the Hausdorff dimension

A natural next step in metric theory of Diophantine approximation was generalisation of the inequalities (1) to $|xq - p| < q^{-s}$, $s > 1$, and $|P(x)| < H^{-w}$, $w > n$.

For $s > 1$ and $w > n$ theorems 4 and 5 yield the equality $\mu \mathcal{K}_n(\psi) = 0$, i. e. the sets $\mathcal{K}_n(\psi)$ are indistinguishable in terms of the Lebesgue measure. This motivated researchers to study the Hausdorff dimension of these sets.

Vojtěch Jarník [43] and Abram Besicovitch [44] independently proved that

$$\dim \mathcal{K}_1(q^{-s}) = \frac{2}{s+1}.$$

In the paper [24] A. Baker and W. Schmidt considered, in addition to the set $\mathcal{K}_n(\psi)$, the set $\mathcal{T}_n(v)$, $v > n + 1$, of real numbers $x \in I$ such that the inequality

$$|x - \alpha| < H^{-v}, \quad v > n + 1,$$

has infinitely many solutions in algebraic numbers α of degree at most n and height at most $H = H(\alpha)$. They introduced the concept of a regular system and proved the following theorem.

Theorem 6. *The following equalities hold*

$$\dim \mathcal{T}_n(v) = \frac{n+1}{v},$$

$$\frac{n+1}{w+1} \leq \dim \mathcal{K}_n(w) \leq 2 \frac{n+1}{w+1}. \quad (11)$$



Theorem 6 was strengthened in [45], where the upper estimate in (11) was replaced by $\frac{n+1}{w+1}$, thus proving the equality

$$\dim \mathcal{K}_n(w) = \frac{n+1}{w+1}.$$

In the paper [46], Yuri Melnichuk obtained estimates for the Hausdorff dimension of the set of points in the unit circle and the unit sphere with a given order of approximation by rational numbers.

Theorem 6 was generalised for the field of p -adic numbers [47]. Later, regular systems have been constructed in the space $\mathbb{R} \times \mathbb{C} \times \mathbb{Q}_p$ from the result [48] and its generalisations [49].

The concept of regular systems of points introduced by A. Baker and W. Schmidt in [24] led to lower bounds for the Hausdorff dimensions of sets of real numbers where an integer polynomial and its derivative provide a given order of approximation of zero. Note that problems related to the Hausdorff measure are closely related to Wirsing's conjecture [50; 51].

A collection of papers by Detta Dickinson, Maurice Dodson, Victor Beresnevich, Vasily Bernik and Sanju Velani [33; 52; 53] introduced the concept of ubiquitous systems, which are in many aspects similar to regular systems. This concept was used to prove lower bounds for the Hausdorff dimension of the sets of real points related to theorem 5.

Interesting facts related to Hausdorff measure of Diophantine sets were obtained by Bryan Rynne [54]. It turns out that in certain cases the Hausdorff dimension is independent of the measure of solutions of Diophantine inequalities.

Let $\bar{\tau} = (\tau_1, \dots, \tau_n) \in \mathbb{R}_+^n$ and $\tau_1 \geq \dots \geq \tau_n$, $\sum_{i=1}^n \tau_i \geq n$.

Let $W_n(\tau) = \{x \in \mathbb{R}^n : |qx_i - p_i| < q^{-\tau_i}, 1 \leq i \leq n, \text{ for infinitely many } (p, q) \in \mathbb{Z}^n \times \mathbb{N}\}$.

Then

$$\dim W_n(\tau) = \min_{1 \leq j \leq n} \frac{n+1 + \sum_{i=j+1}^n (\tau_j - \tau_i)}{\tau_j + 1}.$$

For $n=2$, Rynne's theorem was generalised for rational approximation of the points of the curve $f \in C^{(3)}(I_0)$ defined on an interval I_0 :

$$C_f = \{(x, f(x)) : x \in I_0\}.$$

Let $\bar{\tau} = (\tau_1, \tau_2)$, where τ_1 and τ_2 are positive numbers such that $0 < \min(\tau_1, \tau_2) < 1$ and $\tau_1 + \tau_2 \geq 1$. Assume that

$$\dim \{x \in I_0 : f''(x) = 0\} \leq \frac{2 - \min(\tau_1, \tau_2)}{1 + \max(\tau_1, \tau_2)}.$$

Then

$$\dim W_2(\bar{\tau}) \cap C_f = \frac{2 - \min(\tau_1, \tau_2)}{1 + \max(\tau_1, \tau_2)}.$$

Similar results have been obtained for approximation by real algebraic numbers [15; 55].

Victor Beresnevich and Evgeni Zorin proved the following fact [56].

Let M be a twice continuously differentiable submanifold of \mathbb{R}^n of codimension m , and let

$$\frac{1}{n} \leq \tau \leq \frac{1}{m}.$$

Then

$$\dim S_n(\tau) \cap M \geq s := \frac{n+1}{\tau+1} - m.$$

Furthermore,

$$H^s(S_n(\tau) \cap M) = H^s(M).$$

Recently, Victor Beresnevich, Robert Vaughan, Sanju Velani and Evgeni Zorin [57] worked on finding an upper bound on the quantity of rational points within a ψ -neighbourhood of manifolds. Using this result, they proved the following theorem.



Theorem 7. Let $M_f \subset \mathbb{R}^n$ be a manifold defined on an open subset $U \subset \mathbb{R}^d$, and let

$$H^s \left(\left\{ \alpha \in U : \left| \det \left(\frac{\partial^2 f_j}{\partial \alpha_i \partial \alpha_j} \right)_{1 \leq i, j \leq n} \right| = 0 \right\} \right) = 0$$

for $s = \frac{n+1}{\tau+1} - m$.

If $d > \frac{n+1}{2}$ and $\frac{1}{n} \leq \tau \leq \frac{1}{2m+1}$, then

$$\dim S_n(\tau) \cap M_f \leq s.$$

Generalisations of this result have been obtained by David Simmons and Jing-Jing Huang [58; 59].

Theorem 8. Let $M := \{(x, f(x)) : x \in U \subset \mathbb{R}^d\}$, where $f : U \rightarrow \mathbb{R}^m$, $f \in C^{(2)}$. Let $\bar{\tau} = (\tau_1, \tau_2, \dots, \tau_n) \in \mathbb{R}_{>0}^n$ with

$$\tau_1 \geq \dots \geq \tau_d \geq \max_{d+1 \leq i \leq n} \left\{ \tau_i, \frac{1 - \sum_{j=1}^{n-m} \tau_{j+d}}{d} \right\} \text{ and } \sum_{i=1}^m \tau_{d+i} < 1.$$

Then

$$\dim(W_n(\tau) \cap M) \geq \min_{1 \leq j \leq d} \left\{ \frac{n+1 + \sum_{i=j}^m (\tau_j - \tau_i)}{\tau_j + 1} - m \right\}.$$

Along with the result of V. Beresnevich and E. Zorin, this yields the exact value of $\dim(W_n(\tau) \cap M)$. The proofs are based on the mass transference principle developed by V. Beresnevich and S. Velani [60].

Let us note several other notable results on Hausdorff dimension of sets of numbers with various Diophantine properties. Irina Morotskaya generalised the Baker – Schmidt – Bernik theorem for the field of p -adic numbers [47], and Dickinson and Dodson proved a generalisation of this theorem for non-degenerate curves [53]. In 2017 Victor Beresnevich, Jung-Jo Lee and Robert Vaughan proved a sharp lower bound for a set of τ -approximable numbers in a C^2 submanifold of \mathbb{R}^n for $\frac{1}{n} \leq \tau < \frac{1}{m}$. Rynne’s result was improved in a paper by Konstantin Yavid [61].

Results of this section were strengthened and generalised over the last few years [54; 59; 60; 62–71].

Diophantine approximation in complex and p -adic cases

In Sprindžuk’s monograph [4], Mahler’s problem was generalised to the fields of complex and p -adic numbers. Generalisations of Khintchine’s theorem for complex numbers were obtained by Denis Vasilyev [28]. Diophantine approximation in the field \mathbb{C} was studied in the papers by Irina Morozova [72], Dmitry Kleinbock and George To-manov [73], Natalia Sakovich [74]. An in-depth discussion of simultaneous Diophantine approximation can be found in [62]. The main results of these papers are complete analogues of Khintchine’s theorem and the Jarnik – Besicovitch theorem for weighted simultaneous Diophantine approximation in the p -adic case, as well as a lower bound for the Hausdorff dimension of weighted simultaneously approximable points on p -adic curves [75].

Simultaneous Diophantine approximation

The results that gave rise to metric theory of Diophantine approximation were the Khintchine – Groshev theorem and Mahler’s problem. J. Kubilius obtained a complete solution of Mahler’s problem for $n = 2$ by first considering simultaneous approximation of a parabola by rational points in \mathbb{R}^2 and applying Khintchine’s transference principle [10].

The next major problem in simultaneous approximation was posed by V. Sprindžuk [3; 4].

For a fixed vector $\bar{x} = (x_1, \dots, x_k) \in \mathbb{R}^k$, $1 \leq k \leq n$, let $w \leq w_n(x)$ be the exact upper bound of positive $w_1 > 0$ such that the system of inequalities

$$\max_{1 \leq j \leq k} \left(|P(x_j)| < H^{-w_1} \right)$$



has infinitely many solutions in polynomials $P(t) \in \mathbb{Z}[x]$ of degree $\deg P = n$ and height $H(P) = H$. V. Sprindžuk conjectured that $w = \frac{n-k+1}{k}$, which was proved in [76]. In 1980 V. Sprindžuk [5] posed a problem of approximating points in the space $\Omega = \mathbb{R} \times \mathbb{C} \times \mathbb{Q}_p = \{(x, z, \omega)\}$ by algebraic numbers from \mathbb{R} , \mathbb{C} and \mathbb{Q}_p . He assumed that the following statement was true. Let μ_1 be the Lebesgue measure in \mathbb{R} and \mathbb{C} , and μ_2 be the Haar measure in \mathbb{Q}_p . For almost all $\bar{u} = (x, z, \omega)$ (with respect to the product measure $\mu_1 \times \mu_2$) the system of inequalities

$$|P(x)| < H^{-v_1}, \quad |P(z)| < H^{-v_2}, \quad |P(\omega)| < H^{-v_3},$$

where $v_j \geq -1$, $j = 1, 2$, $v_3 \geq 0$, has only a finite number of solutions in polynomials $P(t) \in \mathbb{Z}[t]$.

This conjecture was proved by Frantz Želudevič [77]. The next major step was generalisation of the Želudevič's results to systems of inequalities where the right-hand sides are arbitrary functions $\psi(H)$. Analogues of Khintchine's theorem were proved for both convergence and divergence cases [78–80].

Of particular interest is the paper [81] that made use of Kleinbock and Margulis's method [49; 68; 70; 71; 76; 78; 79; 82–92].

A number of papers in theory of Diophantine approximations are devoted to studying the distances between conjugate algebraic numbers. The papers [88; 92] can be considered as an introduction to this topic. In the paper [93] Victor Beresnevich, Vasili Bernik and Friedrich Götze applied methods of metric theory of Diophantine approximation to such problems. It was proved that the inequality $|\alpha_1 - \alpha_2| \ll Q^{(n+1)/3}$ not only has a solution in algebraic conjugate numbers α_1 and α_2 of degree n and height $H(\alpha_1) = H(\alpha_2) = Q$, but also the number of the respective minimal polynomials was estimated from below by $c(n)Q^{(n+1)/3}$. This method was generalised for the distribution of polynomial root clusters [93; 94].

Distribution of rational points close to curves and surfaces

In [1; 2] J. Kubilius showed how estimates of the number of rational points close to a parabola $G_2 = (x, x^2)$ lead to a proof of Mahler's conjecture for $n = 2$. The proof was based on Khintchine's transference principle [10]. It turned out that metric theorems concerned with estimation of a dot product between an integer vector $\bar{a} = (a_n, \dots, a_1, a_0)$ and a vector-function $F(x) = (f_n(x), \dots, f_1(x), 1)$ are stronger than metric theorems in simultaneous Diophantine approximation, which is implied by certain results from geometry of numbers and dynamical system theory [35]. In 1994 Martin Huxley [95] proved a theorem on distribution of rational numbers close to smooth

curves, estimating the number $N_f(Q, \delta, J) := \#\left\{\left(\frac{p_1}{q}, \frac{p_2}{q}\right) \in \mathbb{Q}^2 : \frac{p_1}{q} \in J, \left|f\left(\frac{p_1}{q}\right) - \frac{p_2}{q}\right| \leq \delta, 0 < q < Q\right\}$.

Let $I \subset \mathbb{R}$ be a compact interval, c_1 and c_2 be positive constants, and let $F(I; c_1, c_2)$ be the set of functions $f: I \rightarrow \mathbb{R}$, $f \in C^{(2)}$, such that

$$c_1 \leq |f''(x)| \leq c_2 \quad \forall x \in I.$$

M. Huxley proved that

$$N_f(Q, \delta, I) \ll_{\delta} c^{10/3} \delta^{1-\delta} Q^2 + c^{1/3}, \quad c = \max(c_2, c_1^{-1}).$$

For $\delta > Q^{2/3}$, Huxley's result was improved in [52; 55; 56], where it was shown that for any $f \in F(I; c_1, c_2)$, any $Q > 1$ and $0 < \delta < \frac{1}{2}$, we have

$$N_f(Q, \delta, I) \ll \delta Q^2 + \delta^{-1/2} Q.$$

Later the quantity $N_f(Q, \delta, I)$ was estimated from below, and from above and below in the non-homogeneous case [1; 52; 53; 81].

Distribution of algebraic numbers

Distribution of rational numbers in real intervals is quite well-studied and can be described through Farey sequences. However, until recently, not much was known about distribution of algebraic numbers, even if their degrees were small. In 1985 V. Bernik was shown a letter from K. Mahler to V. Sprindžuk, where K. Mahler expressed his surprise about the many unanswered questions related to distribution of algebraic numbers both



on the real line and in the complex plane. It wasn't until the 2010s that Denis Koleda began to solve the problems posed by K. Mahler [96–100]. Let us discuss the main results that he obtained.

Let A_n be the set of real algebraic numbers of degree n , and let the counting function of such algebraic numbers of height at most Q in the interval I be defined as

$$\Phi_n(Q, I) := \#\{\alpha \in A_n \cap I : H(\alpha) \leq Q\}.$$

Theorem 9 [99; 100]. *There exists a continuous positive function $\varphi_n(x)$ such that for any interval $I \subseteq \mathbb{R}$, we have*

$$\Phi_n(Q, I) = \frac{Q^{n+1}}{2\zeta(n+1)} \int_I \varphi_n(x) dx + O\left(Q^n (\ln Q)^{l(n)}\right),$$

where $l(n) = 0$ for $n \geq 3$, $l(n) = 1$ for $n = 2$ and the implicit constant in the big- O notation only depends on n .

The function $\varphi_n(x)$ can be defined explicitly. The remainder term in theorem 9 was shown to be sharp [96], up to a constant, for all n . Further studies of the distribution of algebraic numbers by Denis Koleda, Friedrich Götze and Dmitri Zaporozhets [97] led to the following description of the density of points with algebraically conjugate coordinates in the space $\mathbb{R}^k \times \mathbb{C}^l$.

Let $A_n(k, l)$ be the set of points in the space $\mathbb{R}^k \times \mathbb{C}^l$ such that their coordinates are roots of the same irreducible integral polynomial of degree n (i. e. k real and l complex conjugate algebraic numbers of degree n over \mathbb{Q}). Let us define a function

$$\Phi_{k,l}(Q, B) := \#\{\alpha \in A_n(k, l) \cap B : H(\alpha) \leq Q\}$$

for $Q \geq 1$ and $B \subset \mathbb{R}^k \times \mathbb{C}^l$.

Theorem 10 [97]. *Let $B \subset \mathbb{R}^k \times \mathbb{C}^l$ be a region such that its boundary ∂B is contained in a finite union of Lipschitz transformations of the cube $[0, 1]^{k+2l-1}$. Then*

$$\left| \frac{\Phi_{k,l}(Q, B)}{Q^{n+1}} - \frac{\text{Vol}(B_H)}{2\zeta(n+1)} \int_B \rho_{k,l}(v) dv \right| \leq \begin{cases} cQ^{-1} \log Q, & \text{if } n = 2 \text{ and } l = 0, \\ cQ^{-1} & \text{otherwise,} \end{cases}$$

where $\rho_{k,l} : \mathbb{R}^k \times \mathbb{C}^l \rightarrow \mathbb{R}$ is an explicitly defined continuous non-negative function, dv is a volume element in the space $\mathbb{R}^k \times \mathbb{C}^l$ (considered as \mathbb{R}^{k+2l}), $\zeta(\cdot)$ is the Riemann zeta function, and $\text{Vol}(B_H)$ is the volume of an $(n+1)$ -dimensional region B_H which is the cube $[-1, 1]^{n+1}$ in the case when H is the naive height.

Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be a continuously differentiable function on a finite interval $J \subset \mathbb{R}$, and let $\alpha = (\alpha_1, \alpha_2)$ be a point with algebraically conjugate coordinates such that the minimal polynomial of α_1, α_2 is of degree $\leq n$ and height $\leq Q$. Denote by $M_\varphi^n(Q, \gamma, J)$ the set of points α such that $|\varphi(\alpha_1) - \alpha_2| \leq c_1(n)Q^{-\gamma}$. We show for $0 < \gamma < 1$ and any sufficiently large Q there exist positive values $c_2 < c_3$, where $c_i = c_i(n)$, $i = 2, 3, \dots$, that are independent of Q and the following estimate holds [22; 24; 94; 98–103]

$$c_2 Q^{n+1-\gamma} < \#M_\varphi^n(Q, \gamma, J) < c_3 Q^{n+1-\gamma}.$$

The Mahler – Sprindžuk problem with a non-monotonic right-hand side

It is well known that in the case of convergence, the Khintchine – Groshev theorem also holds for non-monotonic functions $\psi(q)$. A number of researchers have posed other problems in Diophantine approximation with non-monotonic functions in the right-hand side. Generally, the solution then depends on convergence or divergence of the Khintchine-type series with $\psi(q)$ replaced by a different function. For the Mahler – Sprindžuk problem, V. Beresnevich showed in 2005 [27] that the theorem holds for a non-monotonic function $\psi(q)$.

In the papers [104; 105] Natalia Budarina generalised Beresnevich's result for the fields of complex and p -adic numbers, as well as non-degenerate curves. She used Sprindžuk's method, imposing progressively lower bounds on the size of the derivative, and finally applying Kleinbock and Marguli's method for the smallest derivatives [106].



Inhomogeneous Diophantine approximation

The results of the previous sections may be viewed as approximation of zero by values of linear combinations of the form $F(x) = a_n f_n(x) + \dots + a_1 f_1(x) + a_0 \cdot 1$ that are homogeneous with respect to functions $f_n(x), \dots, f_1(x), f_0(x) \equiv 1$. Minkowski's theorems on linear forms and successive minima provide a powerful mechanism for solving this type of problems [10; 14; 15; 19]. However, these methods have very limited utility for derivation of upper and lower bounds if the function $F(x)$ assumes the form $P_1(x) = a_n x^n + \dots + a_1 x + a_0 + \pi$ or $P_2(x) = a_n x^n + \dots + a_1 x + a_0 + \sin x$. These latter problems are known as inhomogeneous Diophantine approximation. A number of metric theorems related to a wide range of functions of the form $P_1(x)$ and $P_2(x)$ have been proved to date [60; 106–110].

Applications of metric theory of Diophantine approximation

The earliest applications of Diophantine approximation to celestial mechanics are probably due to Karl Siegel, who mentions them in his lecture notes. Vladimir Arnold [111] used the Khintchine – Groshev theorem during the development of Kolmogorov – Arnold – Moser theory to prove that almost all celestial systems similar to the Solar system are stable. Systematic application of metric theory of Diophantine approximation in small denominator problems in equations of mathematical physics is described in Ptashnik's monograph [112].

In 2021 a textbook dedicated to applications of metric theory of Diophantine approximation in wireless communications is due to be published by Springer Verlag. Let us point out the articles [31; 113–116] which can be regarded as an introduction to this field of applications, as well as the article by Victor Beresnevich and Sanju Velani, Faustin Adiceam, Jason Levesley and Evgeni Zorin [31] explaining the main ideas behind applications of Diophantine approximation to radio engineering [117; 118].

Biography

Academician J. Kubilius was born in Fermos village, Jurbarkas district of Lithuania. In 1940 he graduated from a grammar school in Raseiniai, in 1946 graduated from Vilnius University, and finished his postgraduate studies at Leningrad University in 1951 under scientific supervision of academician Yuri Linnik. At the age of 36 J. Kubilius successfully defended his doctor of science thesis. Between 1958 and 1992 he was the rector of Vilnius University.

J. Kubilius was the scientific advisor of Belarusian mathematicians Vladimir Sprindžuk (full member of the Academy of Sciences of BSSR) and Nikolai Lazakovich (doctor of science). The academic school of number theory in Belarus, which has 5 doctor of science degree holders and over 40 PhDs degree holders, owes a lot to the work of J. Kubilius. He was the scientific advisor of Ramunė Sliesoraitienė [119], whose PhDs thesis was devoted to metric theory of Diophantine approximation. He was an opponent during thesis defenses of Belarusian mathematicians Vasili Bernik, Ella Kovalevskaya, Vladimir Mashanov.

A full list of scientific degrees and honors of academician J. Kubilius takes up an entire printed page. For more details, we refer the reader to the article [120].

Библиографические ссылки

1. Кубилюс ЙП. О применении метода академика Виноградова к решению одной задачи метрической теории чисел. *Доклады Академии наук СССР*. 1949;67:783–786.
2. Кубилюс ЙП. Об одной метрической проблеме теории диофантовых приближений. *Доклады Академии наук Литовской ССР*. 1959;2:3–7.
3. Спринджук ВГ. Доказательство гипотезы Малера о мере множества S -чисел. *Известия Академии наук СССР. Серия математическая*. 1965;29(2):379–436.
4. Спринджук ВГ. *Проблема Малера в метрической теории чисел*. Минск: Наука и техника; 1967. 181 с.
5. Спринджук ВГ. Достижения и проблемы теории диофантовых приближений. *Успехи математических наук*. 1980;35(4): 3–68.
6. Borel MÈ. Les probabilités dénombrables et leurs applications arithmétiques. *Rendiconti del Circolo Matematico di Palermo (1884–1940)*. 1909;27:247–271. DOI: 10.1007/bf03019651.
7. Khintchine A. Einige Sätze über Kettenbrüche, mit Anwendungen auf die Theorie der diophantischen Approximationen. *Mathematische Annalen*. 1924;92:115–125.
8. Грошев АВ. Теорема о системе линейных форм. *Доклады Академии наук СССР*. 1938;19:151–152.
9. Harman G. *Metric Number Theory*. Oxford: Clarendon Press; 1998. 297 p. (London Mathematical Society monographs; new series 18).
10. Cassels JWS. *An introduction to Diophantine approximation*. 1st edition. Cambridge: Cambridge University Press; 1957. 166 p. (Cambridge tracts in mathematics and mathematical physics; No. 45).



11. Sprindžuk VG. *Mahler's problem in metric number theory*. Volkmann B, translator. Providence: American Mathematical Society; 1969. 192 p. (Translations of mathematical monographs; volume 25).
12. Khintchine A. Zwei Bemerkungen zu einer Arbeit des Herrn Perron. *Mathematische Zeitschrift*. 1925;22:274–284. DOI: 10.1007/bf01479606.
13. Mahler K. Über das Maß der Menge aller S -Zahlen. *Mathematische Annalen*. 1932;106:131–139.
14. Schmidt WM. Bounds for certain sums; a remark on a conjecture of Mahler. *Transactions of the American Mathematical Society*. 1961;101(2):200–210. DOI: 10.1090/s0002-9947-1961-0132036-2.
15. Bugeaud Y. *Approximation by algebraic numbers*. Cambridge: Cambridge University Press; 2004. 290 p. (Cambridge tracts in mathematics; volume 160). DOI: 10.1017/CBO9780511542886.
16. Koksma JF. Über die Mahlersche Klasseneinteilung der transzendenten Zahlen und die Approximation komplexer Zahlen durch algebraische Zahlen. *Monatshefte für Mathematik*. 1939;48(1):176–189. DOI: 10.1007/bf01696176.
17. LeVeque WJ. Note on S -numbers. *Proceedings of the American Mathematical Society*. 1953;4:189–190. DOI: 10.1090/s0002-9939-1953-0054659-2.
18. Kasch F, Volkmann B. Zur Mahlerschen Vermutung über S -Zahlen. *Mathematische Annalen*. 1958;136(5):442–453. DOI: 10.1007/BF01347794.
19. Schmidt WM. Metrische Sätze über simultane Approximation abhängiger Größen. *Monatshefte für Mathematik*. 1964;68(2):154–166. DOI: 10.1007/bf01307118.
20. Volkmann B. The real cubic case of Mahler's conjecture. *Mathematika*. 1961;8(1):55–57. DOI: 10.1112/s0025579300002126.
21. Davenport H. A note on binary cubic forms. *Mathematika*. 1961;8(1):58–62. DOI: 10.1112/s0025579300002138.
22. Bernik V, Götzte F, Gusakova A. On points with algebraically conjugate coordinates close to smooth curves. *Записки научных семинаров ПОМИ*. 2016;448:14–47.
23. Cassels JWS. Some metrical theorems in Diophantine approximation: v. on a conjecture of Mahler. *Mathematical Proceedings of the Cambridge Philosophical Society*. 1951;47(1):18–21. DOI: 10.1017/s0305004100026323.
24. Baker A, Schmidt WM. Diophantine approximation and Hausdorff dimension. *Proceedings of the London Mathematical Society*. 1970;s3-21(1):1–11. DOI: 10.1112/plms/s3-21.1.1.
25. Берник В. О точном порядке приближения нуля значениями целочисленных многочленов. *Acta Arithmetica*. 1989–1990;53(1):17–28.
26. Beresnevich V. On approximation of real numbers by real algebraic numbers. *Acta Arithmetica*. 1999;90(2):97–112. DOI: 10.4064/aa-90-2-97-112.
27. Beresnevich V. On a theorem of V. Bernik in the metric theory of Diophantine approximation. *Acta Arithmetica*. 2005;117(1):71–80. DOI: 10.4064/aa117-1-4.
28. Берник ВИ, Васильев ДВ. Теорема типа Хинчина для целочисленных многочленов комплексной переменной. *Труды Института математики НАН Беларуси*. 1999;3:10–20.
29. Beresnevich VV, Bernik VI, Kovalevskaya EI. On approximation of p -adic numbers by p -adic algebraic numbers. *Journal of Number Theory*. 2005;111(1):33–56. DOI: 10.1016/j.jnt.2004.09.007.
30. Mohammadi A, Gosefidy AS. S -arithmetic Khintchine-type theorem. *Geometric and Functional Analysis*. 2009;19(4):1147–1170. DOI: 10.1007/s00039-009-0029-z.
31. Adiceam F, Beresnevich V, Levesley J, Velani S, Zorin E. Diophantine approximation and applications in interference alignment. *Advances in Mathematics*. 2016;302:231–279. DOI: 10.1016/j.aim.2016.07.002.
32. Берник ВИ, Шамукова НВ. Приближение действительных чисел целыми алгебраическими числами и теорема Хинчина. *Доклады Национальной академии наук Беларуси*. 2006;50(3):30–32.
33. Bernik VI, Dodson MM. *Metric Diophantine approximation on manifolds*. Cambridge: Cambridge University Press; 1999. 172 p. (Cambridge tracts in mathematics; volume 137). DOI: 10.1017/CBO9780511565991.
34. Beresnevich V, Bernik V. On a metrical theorem of W. Schmidt. *Acta Arithmetica*. 1996;75(3):219–233. DOI: 10.4064/aa-75-3-219-233.
35. Kleinbock DY, Margulis GA. Flows on homogeneous spaces and Diophantine approximation on manifolds. *Annals of Mathematics*. 1998;148(1):339–360. DOI: 10.2307/120997.
36. Beresnevich V. A Groshev type theorem for convergence on manifolds. *Acta Mathematica Hungarica*. 2002;94(1–2):99–130. DOI: 10.1023/A:1015662722298.
37. Bernik V, Kleinbock D, Margulis G. Khintchine-type theorems on manifolds: the convergence case for standard and multiplicative versions. *International Mathematics Research Notices*. 2001;2001(9):453–486. DOI: 10.1155/S1073792801000241.
38. Beresnevich VV, Bernik VI, Kleinbock DYa, Margulis GA. Metric Diophantine approximation: the Khintchine – Groshev theorem for non-degenerate manifolds. *Moscow Mathematical Journal*. 2002;2(2):203–225. DOI: 10.17323/1609-4514-2002-2-2-203-225.
39. Baker A. On a theorem of Sprindžuk. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*. 1966;292(1428):92–104. DOI: 10.1098/rspa.1966.0121.
40. Koksma JF. *Diophantische approximationen*. Berlin: Springer; 1974. 172 p. (Ergebnisse der Mathematik und ihrer Grenzgebiete; volume 4). DOI: 10.1007/978-3-642-65618-7.
41. Allen D, Beresnevich V. A mass transference principle for systems of linear forms and its applications. *Compositio Mathematica*. 2018;154(5):1014–1047. DOI: 10.1112/s0010437x18007121.
42. Берник ВИ, Васильев ДВ, Засимович ЕВ. Диофантовы приближения с постоянной правой частью неравенств на коротких интервалах. *Доклады Национальной академии наук Беларуси*. 2021;65(4):397–403. DOI: 10.29235/1561-8323-2021-65-4-397-403.
43. Jarnik V. Diophantische approximationen und Hausdorffsches mass. *Математический сборник*. 1929;36(3–4):371–382.
44. Besicovitch AS. Sets of fractional dimensions (IV): on rational approximation to real numbers. *Journal of the London Mathematical Society*. 1934;s1-9(2):126–131. DOI: 10.1112/jlms/s1-9.2.126.
45. Берник ВИ. Применение размерности Хаусдорфа в теории диофантовых приближений. *Acta Arithmetica*. 1983;42:219–253. DOI: 10.4064/aa-42-3-219-253.
46. Мельничук ЮВ. Диофантовы приближения на окружности и размерность Хаусдорфа. *Математические заметки*. 1979;26(3):347–354.



47. Берник ВИ, Моротская ИЛ. Диофантовы приближения в \mathbb{Q}_p и размерность Хаусдорфа. *Весті Нацыянальнай акадэміі навук Беларусі. Серыя фізіка-матэматычных навук*. 1986;3:3–9.
48. Берник ВИ, Калоша НИ. Приближение нуля значениями целочисленных полиномов в пространстве $\mathbb{R} \times \mathbb{C} \times \mathbb{Q}_p$. *Весті Нацыянальнай акадэміі навук Беларусі. Серыя фізіка-матэматычных навук*. 2004;1:121–123.
49. Бударина НВ. Метрическая теория совместных диофантовых приближений в $\mathbb{R}^k \times \mathbb{C}^l \times \mathbb{Q}_p^m$. *Чебышевский сборник. посвящается 65-й годовщине со дня рождения профессора Сергея Михайловича Воронина*. 2011;12(1):17–50.
50. Badziahin D, Schleischitz J. An improved bound in Wirsing's problem. *Transactions of the American Mathematical Society*. 2021;374:1847–1861. DOI: 10.1090/tran/8245.
51. Берник ВИ, Тищенко КИ. Целочисленные многочлены с перепадами высот коэффициентов и гипотеза Вирзинга. *Доклады Национальной академии наук Беларуси*. 1993;37(5):9–11.
52. Beresnevich V, Dickinson D, Velani S. Diophantine approximation on planar curves and the distribution of rational points (with an Appendix by R. C. Vaughan). *Annals of Mathematics*. 2007;166(2):367–426. DOI: 10.4007/annals.2007.166.367.
53. Dickinson H, Dodson MM. Extremal manifolds and Hausdorff dimension. *Duke Mathematical Journal*. 2000;101(2):271–281. DOI: 10.1215/S0012-7094-00-10126-3.
54. Rynne BP. Simultaneous Diophantine approximation on manifolds and Hausdorff dimension. *Journal of Number Theory*. 2003;98(1):1–9. DOI: 10.1016/s0022-314x(02)00035-5.
55. Кудин АС, Луневич АВ. Аналог теоремы Хинчина в случае расходимости в полях действительных, комплексных и p -адических чисел. *Труды Института математики*. 2015;23(1):76–83.
56. Beresnevich V, Zorin E. Explicit bounds for rational points near planar curves and metric Diophantine approximation. *Advances in Mathematics*. 2010;225(6):3064–3087. DOI: 10.1016/j.aim.2010.05.021.
57. Beresnevich V, Vaughan RC, Velani S, Zorin E. Diophantine approximation on manifolds and the distribution of rational points: contributions to the convergence theory. *International Mathematics Research Notices*. 2017;2017(10):2885–2908. DOI: 10.1093/imrn/rnv389.
58. Huang J-J. The density of rational points near hypersurfaces. *Duke Mathematical Journal*. 2020;169(11):2045–2077. DOI: 10.1215/00127094-2020-0004.
59. Simmons D. Some manifolds of Khinchin type for convergence. *Journal de Théorie des Nombres de Bordeaux*. 2018;30(1):175–193. DOI: 10.5802/jtnb.1021.
60. Beresnevich V, Velani S. An inhomogeneous transference principle and Diophantine approximation. *Proceedings of the London Mathematical Society*. 2010;101(3):821–851. DOI: 10.1112/plms/pdq002.
61. Явид КЮ. Оценка размерности Хаусдорфа множеств сингулярных векторов. *Доклады Академии наук БССР*. 1987;31(9):777–780.
62. Beresnevich V, Levesley J, Ward B. A lower bound for the Hausdorff dimension of the set of weighted simultaneously approximable points over manifolds. *International Journal of Number Theory*. 2021;17(8):1795–1814. DOI: 10.1142/S1793042121500639.
63. Bernik VI. Applications of measure theory and Hausdorff dimension to the theory of Diophantine approximation. *New advances in transcendence theory*. 1988:25–36. DOI: 10.1017/CBO9780511897184.003.
64. Берник ВИ. Применение размерности Хаусдорфа в теории диофантовых приближений. *Acta Arithmetica*. 1983;42:219–253.
65. Beresnevich V, Lee L, Vaughan RC, Velani S. Diophantine approximation on manifolds and lower bounds for Hausdorff dimension. *Mathematika*. 2017;63(3):762–779. DOI: 10.1112/s0025579317000171.
66. Bernik VI, Pereverseva NA. The method of trigonometric sums and lower estimates of Hausdorff dimension. *Analytic and Probabilistic Methods in Number Theory*. 1992;2:75–81. DOI: 10.1515/9783112314234-011.
67. Bugeaud Y. Approximation by algebraic integers and Hausdorff dimension. *Journal of the London Mathematical Society*. 2002;65(3):547–559. DOI: 10.1112/S0024610702003137.
68. Beresnevich VV, Velani SL. A note on simultaneous Diophantine approximation on planar curves. *Mathematische Annalen*. 2007;337(4):769–796. DOI: 10.1007/s00208-006-0055-1.
69. Бересневич ВВ. Применение понятия регулярных систем точек в метрической теории чисел. *Весті Нацыянальнай акадэміі навук Беларусі. Серыя фізіка-матэматычных навук*. 2000;1:35–39.
70. Бересневич ВВ. О построении регулярных систем точек с вещественными, комплексными и p -адическими алгебраическими координатами. *Весті Нацыянальнай акадэміі навук Беларусі. Серыя фізіка-матэматычных навук*. 2003;1:22–27.
71. Kovalevskaya EI, Bernik V. Simultaneous inhomogeneous Diophantine approximation of the values of integral polynomials with respect to Archimedean and non-Archimedean valuations. *Acta Mathematica Universitatis Ostraviensis*. 2006;14(1):37–42.
72. Бернік ВІ, Марозава ІМ. Гіпотэза Бэйкера і рэгулярныя сістэмы алгебраічных лічбаў з абмежаваннем на значэнне вытворнай. *Весті Нацыянальнай акадэміі навук Беларусі. Серыя фізіка-матэматычных навук*. 1996;3:109–113.
73. Kleinbock D, Tomanov G. Flows on S -arithmetic homogeneous spaces and applications to metric Diophantine approximation. *Commentarii Mathematici Helvetici*. 2007;82(3):519–581. DOI: 10.4171/cmh/102.
74. Берник ВИ, Сакович НВ. Регулярные системы комплексных алгебраических чисел. *Доклады Академии наук Беларуси*. 1994;38(5):10–13.
75. Бересневич ВВ, Ковалевская ЭИ. О диофантовых приближениях зависимых величин в p -адическом случае. *Математические заметки*. 2003;25(1):22–37. DOI: 10.4213/mzm165.
76. Берник ВИ. Метрическая теорема о совместном приближении нуля значениями целочисленных многочленов. *Известия Академии наук СССР. Серия математическая*. 1980;44(1):24–45.
77. Želudevich F. Simultane diophantische Approximationen abhängiger Größen in mehreren Metriken. *Acta Arithmetica*. 1986;46(3):285–296. DOI: 10.4064/aa-46-3-285-296.
78. Bernik V, Budarina N, Dickinson D. A divergent Khintchine theorem in the real, complex, and p -adic fields. *Lithuanian Mathematical Journal*. 2008;48(2):158–173. DOI: 10.1007/s10986-008-9005-9.
79. Budarina N, Dickinson D, Bernik V. Simultaneous Diophantine approximation in the real, complex and p -adic fields. *Mathematical Proceedings of the Cambridge Philosophical Society*. 2010;149(2):193–216. DOI: 10.1017/s0305004110000162.
80. Домбровский ИР. Совместные приближения действительных чисел алгебраическими числами ограниченной степени. *Доклады Академии наук БССР*. 1989;33(3):205–208.
81. Beresnevich V. Rational points near manifolds and metric Diophantine approximation. *Annals of Mathematics*. 2012;175(1):187–235. DOI: 10.4007/annals.2012.175.1.5.



82. Beresnevich VV, Velani SL. Simultaneous inhomogeneous Diophantine approximation on manifolds. *Journal of Mathematical Sciences*. 2012;180(5):531–541. DOI: 10.1007/s10958-012-0651-4.
83. Budarina N, Dickinson D, Levesley J. Simultaneous Diophantine approximation on polynomial curves. *Mathematika*. 2010; 56(1):77–85. DOI: 10.1112/s0025579309000382.
84. Budarina N. On a problem of Bernik, Kleinbock and Margulis. *Glasgow Mathematical Journal*. 2011;53(3):669–681. DOI: 10.1017/s0017089511000255.
85. Budarina N, Dickinson D. Simultaneous Diophantine approximation in two metrics and the distance between conjugate algebraic numbers in $\mathbb{R} \times \mathbb{Q}_p$. *Indagationes Mathematicae*. 2012;23(1–2):32–41. DOI: 10.1016/j.indag.2011.09.012.
86. Берник ВИ, Борбат ВН. Совместная аппроксимация нуля значениями целочисленных полиномов. *Труды Математического института имени В. А. Стеклова*. 1997;218:58–73.
87. Bernik VI, Budarina N, O'Donnell H. On regular systems of real algebraic numbers of third degree in short intervals. *Современные проблемы математики*. 2013;17:61–75. DOI: 10.4213/spm43.
88. Bugeaud Y, Mignotte M. Polynômes à coefficients entiers prenant des valeurs positives aux points réels. *Bulletin mathématique de la Société des Sciences Mathématiques de Roumanie*. 2010;53(3):219–224.
89. Bernik V, Götze F. A new connection between metric theory of Diophantine approximations and distribution of algebraic numbers. *Contemporary Mathematics*. 2015;631:33–45. DOI: 10.1090/conm/631/12594.
90. Бударина НВ. Совместные диофантовы приближения с немонотонными правыми частями. *Доклады Академии наук*. 2011;437(4):441–443.
91. Bernik V, Mc Guire S. How small can polynomials be in an interval of given length? *Glasgow Mathematical Journal*. 2020; 62(2):261–280. DOI: 10.1017/S0017089519000077.
92. Bugeaud Y, Mignotte M. Polynomial root separation. *International Journal of Number Theory*. 2010;6(3):587–602. DOI: 10.1142/ s1793042110003083.
93. Beresnevich V, Bernik V, Götze F. The distribution of close conjugate algebraic numbers. *Compositio Mathematica*. 2010; 146(5):1165–1179. DOI: 10.1112/S0010437X10004860.
94. Bernik V, Budarina N, O'Donnell H. Discriminants of polynomials in the Archimedean and non-Archimedean metrics. *Acta Mathematica Hungarica*. 2018;154(2):265–278. DOI: <https://doi.org/10.1007/s10474-018-0794-y>.
95. Huxley MN. The rational points close to a curve. *Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Série 4*. 1994;21(3):357–375.
96. Коледа ДВ. Об асимптотике распределения алгебраических чисел при возрастании их высот. *Чебышевский сборник*. 2015;16(1):191–204. DOI: 10.22405/2226-8383-2015-16-1-191-204.
97. Götze F, Koleda D, Zaporozhets D. Joint distribution of conjugate algebraic numbers: a random polynomial approach. *Advances in Mathematics*. 2020;359:106849. DOI: 10.1016/j.aim.2019.106849.
98. Лебедь ВВ, Берник ВИ. Алгебраические точки на плоскости. *Фундаментальная и прикладная математика*. 2005; 11(6):73–80.
99. Коледа ДВ. О распределении вещественных алгебраических чисел второй степени. *Весті Національної академії наук України. Серія фізико-математичних наук*. 2013;3:54–63.
100. Koleda D. On the density function of the distribution of real algebraic numbers. *Journal de Théorie des Nombres de Bordeaux*. 2017;29(1):179–200. DOI: 10.5802/jtnb.975.
101. Берник ВИ, Гётце Ф, Калоша НИ. О количестве алгебраических чисел в коротких интервалах, содержащих рациональные точки. *Журнал Белорусского государственного университета. Математика. Информатика*. 2019;1:4–11. DOI: 10.33581/ 2520-6508-2019-1-4-11.
102. Бударина НВ, Диккинсон Д, Берник ВИ. Оценки снизу для количества векторов с алгебраическими координатами вблизи гладких поверхностей. *Доклады Национальной академии наук Беларуси*. 2020;64(1):7–12. DOI: 10.29235/1561-8323- 2020-64-1-7-12.
103. Bernik VI, Götze F. Distribution of real algebraic numbers of arbitrary degree in short intervals. *Izvestiya: Mathematics*. 2015;79(1):18–39. DOI: 10.1070/im2015v079n01abeh002732.
104. Budarina N, Dickinson D. Diophantine approximation on non-degenerate curves with non-monotonic error function. *Bulletin of the London Mathematical Society*. 2009;41(1):137–146. DOI: 10.1112/blms/bdn116.
105. Budarina N. Diophantine approximation on the curves with non-monotonic error function in the p -adic case. *Чебышевский сборник*. 2010;11(1):74–80.
106. Badziahin D, Beresnevich V, Velani S. Inhomogeneous theory of dual Diophantine approximation on manifolds. *Advances in Mathematics*. 2013;232(1):1–35. DOI: 10.1016/j.aim.2012.09.022.
107. Badziahin D. Inhomogeneous Diophantine approximation on curves and Hausdorff dimension. *Advances in Mathematics*. 2010;223(1):329–351. DOI: 10.1016/j.aim.2009.08.005.
108. Bernik V, Dickinson H, Yuan J. Inhomogeneous Diophantine approximation on polynomials in \mathbb{Q}_p . *Acta Arithmetica*. 1999; 90(1):37–48. DOI: 10.4064/aa-90-1-37-48.
109. Beresnevich V, Ganguly A, Ghosh A, Velani S. Inhomogeneous dual Diophantine approximation on affine subspaces. *International Mathematics Research Notices*. 2020;12:3582–3613. DOI: 10.1093/imrn/rny124.
110. Beresnevich VV, Vaughan RC, Velani SL. Inhomogeneous Diophantine approximation on planar curves. *Mathematische Annalen*. 2011;349(4):929–942. DOI: 10.1007/s00208-010-0548-9.
111. Арнольд ВИ. Малые знаменатели и проблемы устойчивости движения в классической и небесной механике. *Успехи математических наук*. 1963;18(6):91–192.
112. Пташник БИ. *Некорректные граничные задачи для дифференциальных уравнений с частными производными*. Киев: Наукова думка; 1984. 264 с.
113. Cadambe VR, Jafar SA. Interference alignment and degrees of freedom of the K -user interference channel. *IEEE Transactions on Information Theory*. 2008;54(8):3425–3441. DOI: 10.1109/TIT.2008.926344.
114. Jafar SA. Interference alignment: a new look at signal dimensions in a communication network. *Foundations and Trends in Communications and Information Theory*. 2011;7(1):1–134. DOI: 10.1561/01000000047.
115. Jafar SA, Shamai Sh. Degrees of freedom region of the MIMO X channel. *IEEE Transactions on Information Theory*. 2008; 54(1):151–170. DOI: 10.1109/tit.2007.911262.



116. Beresnevich V, Velani S. Number theory meets wireless communications: an introduction for dummies like us. In: Beresnevich V, Burr A, Nazer B, Velani S, editors. *Number theory meets wireless communications*. [S. l.]: Springer International Publishing; 2020. p. 1–67. DOI: 10.1007/978-3-030-61303-7_1.
117. Budarina N, O'Donnell H. On a problem of Nesterenko: when is the closest root of a polynomial a real number? *International Journal of Number Theory*. 2012;8(3):801–811. DOI: 10.1142/s1793042112500455.
118. Motahari AS, Oveis-Gharan S, Maddah-Ali MA, Khandani AK. Real interference alignment: exploiting the potential of single antenna systems. *IEEE Transactions on Information Theory*. 2014;60(8):4799–4810. DOI: 10.1109/tit.2014.2329865.
119. Слесорайтене Р. Теорема Малера – Спринджука для полиномов третьей степени от двух переменных. (II). *Литовский математический сборник*. 1970;10:791–814.
120. Manstavičius E, Jonas Kubilius 1921–2011. *Lithuanian Mathematical Journal*. 2021;61:285–288. DOI: 10.1007/s10986-021-09522-z.

References

- Kubilius JP. [On application of Academician Vinogradov's method to solving a certain problem in metric number theory]. *Doklady Akademii nauk SSSR*. 1949;67:783–786. Russian.
- Kubilius JP. [On a metrical problem in Diophantine approximation theory]. *Doklady Akademii nauk Litovskoi SSR*. 1959;2:3–7. Russian.
- Sprindžuk VG. [Proof of Mahler's conjecture on the measure of the set of S -numbers]. *Izvestiya Akademii nauk SSSR. Seriya matematicheskaya*. 1965;29(2):379–436. Russian.
- Sprindžuk VG. *Problema Malera v metricheskoj teorii chisel* [Mahler's problem in metric number theory]. Minsk: Nauka i tekhnika; 1967. 181 p. Russian.
- Sprindžuk VG. [Advances and problems in the theory of Diophantine approximation]. *Uspekhi matematicheskikh nauk*. 1980;35(4):3–68. Russian.
- Borel MÈ. Les probabilités dénombrables et leurs applications arithmétiques. *Rendiconti del Circolo Matematico di Palermo (1884–1940)*. 1909;27:247–271. DOI: 10.1007/bf03019651.
- Khintchine A. Einige Sätze über Kettenbrüche, mit Anwendungen auf die Theorie der diophantischen Approximationen. *Mathematische Annalen*. 1924;92:115–125.
- Groshev AV. [Theorem on a system of linear forms]. *Doklady Akademii nauk SSSR*. 1938;19:151–152. Russian.
- Harman G. *Metric number theory*. Oxford: Clarendon Press; 1998. 297 p. (London Mathematical Society monographs; new series 18).
- Cassels JWS. *An introduction to Diophantine approximation*. 1st edition. Cambridge: Cambridge University Press; 1957. 166 p. (Cambridge tracts in mathematics and mathematical physics; No. 45).
- Sprindžuk VG. *Mahler's problem in metric number theory*. Volkman B, translator. Providence: American Mathematical Society; 1969. 192 p. (Translations of mathematical monographs; volume 25).
- Khintchine A. Zwei Bemerkungen zu einer Arbeit des Herrn Perron. *Mathematische Zeitschrift*. 1925;22:274–284. DOI: 10.1007/bf01479606.
- Mahler K. Über das Maß der Menge aller S -Zahlen. *Mathematische Annalen*. 1932;106:131–139.
- Schmidt WM. Bounds for certain sums; a remark on a conjecture of Mahler. *Transactions of the American Mathematical Society*. 1961;101(2):200–210. DOI: 10.1090/s0002-9947-1961-0132036-2.
- Bugeaud Y. *Approximation by algebraic numbers*. Cambridge: Cambridge University Press; 2004. 290 p. (Cambridge tracts in mathematics; volume 160). DOI: 10.1017/CBO9780511542886.
- Koksma JF. Über die Mahlersche Klasseneinteilung der transzendenten Zahlen und die Approximation komplexer Zahlen durch algebraische Zahlen. *Monatshefte für Mathematik*. 1939;48(1):176–189. DOI: 10.1007/bf01696176.
- LeVeque WJ. Note on S -numbers. *Proceedings of the American Mathematical Society*. 1953;4:189–190. DOI: 10.1090/s0002-9939-1953-0054659-2.
- Kasch F, Volkman B. Zur Mahlerschen Vermutung über S -Zahlen. *Mathematische Annalen*. 1958;136(5):442–453. DOI: 10.1007/BF01347794.
- Schmidt WM. Metrische Sätze über simultane Approximation abhängiger Größen. *Monatshefte für Mathematik*. 1964;68(2):154–166. DOI: 10.1007/bf01307118.
- Volkman B. The real cubic case of Mahler's conjecture. *Mathematika*. 1961;8(1):55–57. DOI: 10.1112/s0025579300002126.
- Davenport H. A note on binary cubic forms. *Mathematika*. 1961;8(1):58–62. DOI: 10.1112/s0025579300002138.
- Bernik V, Götze F, Gusakova A. On points with algebraically conjugate coordinates close to smooth curves. *Zapiski nauchnykh seminarov POMI*. 2016;448:14–47.
- Cassels JWS. Some metrical theorems in Diophantine approximation: v. on a conjecture of Mahler. *Mathematical Proceedings of the Cambridge Philosophical Society*. 1951;47(1):18–21. DOI: 10.1017/s0305004100026323.
- Baker A, Schmidt WM. Diophantine approximation and Hausdorff dimension. *Proceedings of the London Mathematical Society*. 1970;s3-21(1):1–11. DOI: 10.1112/plms/s3-21.1.1.
- Bernik V. [On the exact order of approximation of zero by values of integer polynomials]. *Acta Arithmetica*. 1989–1990;53(1):17–28. Russian.
- Beresnevich V. On approximation of real numbers by real algebraic numbers. *Acta Arithmetica*. 1999;90(2):97–112. DOI: 10.4064/aa-90-2-97-112.
- Beresnevich V. On a theorem of V. Bernik in the metric theory of Diophantine approximation. *Acta Arithmetica*. 2005;117(1):71–80. DOI: 10.4064/aa117-1-4.
- Bernik VI, Vasilyev DV. [A Khintchine-type theorem for integer polynomials with a complex variable]. *Trudy Instituta matematiki NAN Belarusi*. 1999;3:10–20. Russian.
- Beresnevich VV, Bernik VI, Kovalevskaya EI. On approximation of p -adic numbers by p -adic algebraic numbers. *Journal of Number Theory*. 2005;111(1):33–56. DOI: 10.1016/j.jnt.2004.09.007.



30. Mohammadi A, Golsefidy AS. S-arithmetic Khintchine-type theorem. *Geometric and Functional Analysis*. 2009;19(4):1147–1170. DOI: 10.1007/s00039-009-0029-z.
31. Adiceam F, Beresnevich V, Levesley J, Velani S, Zorin E. Diophantine approximation and applications in interference alignment. *Advances in Mathematics*. 2016;302:231–279. DOI: 10.1016/j.aim.2016.07.002.
32. Bernik VI, Shamukova NV. [Approximation of real numbers by integer algebraic numbers, and the Khintchine’s theorem]. *Doklady of the National Academy of Sciences of Belarus*. 2006;50(3):30–32. Russian.
33. Bernik VI, Dodson MM. *Metric Diophantine approximation on manifolds*. Cambridge: Cambridge University Press; 1999. 172 p. (Cambridge tracts in mathematics; volume 137). DOI: 10.1017/CBO9780511565991.
34. Beresnevich V, Bernik V. On a metrical theorem of W. Schmidt. *Acta Arithmetica*. 1996;75(3):219–233. DOI: 10.4064/aa-75-3-219-233.
35. Kleinbock DY, Margulis GA. Flows on homogeneous spaces and Diophantine approximation on manifolds. *Annals of Mathematics*. 1998;148(1):339–360. DOI: 10.2307/120997.
36. Beresnevich V. A Groshev type theorem for convergence on manifolds. *Acta Mathematica Hungarica*. 2002;94(1–2):99–130. DOI: 10.1023/A:1015662722298.
37. Bernik V, Kleinbock D, Margulis G. Khintchine-type theorems on manifolds: the convergence case for standard and multiplicative versions. *International Mathematics Research Notices*. 2001;2001(9):453–486. DOI: 10.1155/S1073792801000241.
38. Beresnevich VV, Bernik VI, Kleinbock DY, Margulis GA. Metric Diophantine approximation: the Khintchine – Groshev theorem for non-degenerate manifolds. *Moscow Mathematical Journal*. 2002;2(2):203–225. DOI: 10.17323/1609-4514-2002-2-2-203-225.
39. Baker A. On a theorem of Sprindžuk. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*. 1966;292(1428):92–104. DOI: 10.1098/rspa.1966.0121.
40. Koksma JF. *Diophantische approximationen*. Berlin: Springer; 1974. 172 p. (Ergebnisse der Mathematik und ihrer Grenzgebiete; volume 4). DOI: 10.1007/978-3-642-65618-7.
41. Allen D, Beresnevich V. A mass transference principle for systems of linear forms and its applications. *Compositio Mathematica*. 2018;154(5):1014–1047. DOI: 10.1112/s0010437x18007121.
42. Bernik VI, Vasilyev DV, Zasimovich EV. Diophantine approximation with the constant right-hand side of inequalities on short intervals. *Doklady of the National Academy of Sciences of Belarus*. 2021;65(4):397–403. Russian. DOI: 10.29235/1561-8323-2021-65-4-397-403.
43. Jarnik V. Diophantische approximationen und Hausdorffsches mass. *Matematicheskii sbornik*. 1929;36(3–4):371–382.
44. Besicovitch AS. Sets of fractional dimensions (IV): on rational approximation to real numbers. *Journal of the London Mathematical Society*. 1934;s1-9(2):126–131. DOI: 10.1112/jlms/s1-9-2.126.
45. Bernik VI. [Application of the Hausdorff dimension in the theory of Diophantine approximation]. *Acta Arithmetica*. 1983;42:219–253. Russian. DOI: 10.4064/aa-42-3-219-253.
46. Melnichuk YuV. [Diophantine approximation on a circle and the Hausdorff dimension]. *Matematicheskie zametki*. 1979;26(3):347–354. Russian.
47. Bernik VI, Morotskaya IL. [Diophantine approximation in \mathbb{Q}_p and the Hausdorff dimension]. *Proceedings of the National Academy of Sciences of Belarus. Physics and Mathematics Series*. 1986;3:3–9. Russian.
48. Bernik VI, Kalosha NI. Approximation of zero by values of integer polynomials in the space $\mathbb{R} \times \mathbb{C} \times \mathbb{Q}_p$. *Proceedings of the National Academy of Sciences of Belarus. Physics and Mathematics Series*. 2004;1:121–123. Russian.
49. Budarina NV. [Metric theory of simultaneous Diophantine approximation in $\mathbb{R}^k \times \mathbb{C}^l \times \mathbb{Q}_p^m$]. *Chebyshevskii sbornik. Posvyashchaetsya 65-i godovshchine so dnya rozhdeniya professora Sergeya Mikhailovicha Voronina*. 2011;12(1):17–50. Russian.
50. Badziahin D, Schleisnitz J. An improved bound in Wirsing’s problem. *Transactions of the American Mathematical Society*. 2021;374:1847–1861. DOI: 10.1090/tran/8245.
51. Bernik VI, Tishchenko KI. [Integral polynomials with an overfall of the coefficient values and Wirsing’s theorem]. *Doklady of the National Academy of Belarus*. 1993;37(5):9–11. Russian.
52. Beresnevich V, Dickinson D, Velani S. Diophantine approximation on planar curves and the distribution of rational points (with an Appendix by R. C. Vaughan). *Annals of Mathematics*. 2007;166(2):367–426. DOI: 10.4007/annals.2007.166.367.
53. Dickinson H, Dodson MM. Extremal manifolds and Hausdorff dimension. *Duke Mathematical Journal*. 2000;101(2):271–281. DOI: 10.1215/S0012-7094-00-10126-3.
54. Rynne BP. Simultaneous Diophantine approximation on manifolds and Hausdorff dimension. *Journal of Number Theory*. 2003;98(1):1–9. DOI: 10.1016/s0022-314x(02)00035-5.
55. Kudin AS, Lunevich AV. [An analogue of Khintchine’s theorem in the case of divergence in the fields of real, complex and p -adic numbers]. *Trudy Instituta matematiki*. 2015;23(1):76–83. Russian.
56. Beresnevich V, Zorin E. Explicit bounds for rational points near planar curves and metric Diophantine approximation. *Advances in Mathematics*. 2010;225(6):3064–3087. DOI: 10.1016/j.aim.2010.05.021.
57. Beresnevich V, Vaughan RC, Velani S, Zorin E. Diophantine approximation on manifolds and the distribution of rational points: contributions to the convergence theory. *International Mathematics Research Notices*. 2017;2017(10):2885–2908. DOI: 10.1093/imrn/rnv389.
58. Huang J-J. The density of rational points near hypersurfaces. *Duke Mathematical Journal*. 2020;169(11):2045–2077. DOI: 10.1215/00127094-2020-0004.
59. Simmons D. Some manifolds of Khinchin type for convergence. *Journal de Théorie des Nombres de Bordeaux*. 2018;30(1):175–193. DOI: 10.5802/jtnb.1021.
60. Beresnevich V, Velani S. An inhomogeneous transference principle and Diophantine approximation. *Proceedings of the London Mathematical Society*. 2010;101(3):821–851. DOI: 10.1112/plms/pdq002.
61. Yavid KYu. [An estimate for the Hausdorff dimension of sets of singular vectors]. *Doklady Akademii nauk BSSR*. 1987;31(9):777–780. Russian.
62. Beresnevich V, Levesley J, Ward B. A lower bound for the Hausdorff dimension of the set of weighted simultaneously approximable points over manifolds. *International Journal of Number Theory*. 2021;17(8):1795–1814. DOI: 10.1142/S1793042121500639.
63. Bernik VI. Applications of measure theory and Hausdorff dimension to the theory of Diophantine approximation. *New advances in transcendence theory*. 1988:25–36. DOI: 10.1017/CBO9780511897184.003.



64. Bernik VI. [Application of Hausdorff dimension in the theory of Diophantine approximations]. *Acta Arithmetica*. 1983;42: 219–253. Russian.
65. Beresnevich V, Lee L, Vaughan RC, Velani S. Diophantine approximation on manifolds and lower bounds for Hausdorff dimension. *Mathematika*. 2017;63(3):762–779. DOI: 10.1112/s0025579317000171.
66. Bernik VI, Pereverseva NA. The method of trigonometric sums and lower estimates of Hausdorff dimension. *Analytic and Probabilistic Methods in Number Theory*. 1992;2:75–81. DOI: 10.1515/9783112314234-011.
67. Bugeaud Y. Approximation by algebraic integers and Hausdorff dimension. *Journal of the London Mathematical Society*. 2002; 65(3):547–559. DOI: 10.1112/S0024610702003137.
68. Beresnevich VV, Velani SL. A note on simultaneous Diophantine approximation on planar curves. *Mathematische Annalen*. 2007;337(4):769–796. DOI: 10.1007/s00208-006-0055-1.
69. Beresnevich VV. Application of the concept of regular systems in the Metric theory of numbers. *Proceedings of the National Academy of Sciences of Belarus. Physics and Mathematics Series*. 2000;1:35–39. Russian.
70. Beresnevich VV. [On construction of regular systems of points with real, complex and p -adic algebraic coordinates]. *Proceedings of the National Academy of Sciences of Belarus. Physics and Mathematics Series*. 2003;1:22–27. Russian.
71. Kovalevskaya EI, Bernik V. Simultaneous inhomogeneous Diophantine approximation of the values of integral polynomials with respect to Archimedean and non-Archimedean valuations. *Acta Mathematica Universitatis Ostraviensis*. 2006;14(1):37–42.
72. Bernik VI, Marozava IM. Baker’s conjecture and regular sets of algebraic numbers with a restriction on the value of the derivative. *Proceedings of the National Academy of Sciences of Belarus. Physics and Mathematics Series*. 1996;3:109–113. Belarusian.
73. Kleinbock D, Tomanov G. Flows on S -arithmetic homogeneous spaces and applications to metric Diophantine approximation. *Commentarii Mathematici Helvetici*. 2007;82(3):519–581. DOI: 10.4171/cmh/102.
74. Bernik VI, Sakovich NV. Regular systems of complex algebraic numbers. *Doklady Akademii nauk Belarusi*. 1994;38(5):10–13. Russian.
75. Beresnevich VV, Kovalevskaya EI. [On Diophantine approximation of dependent variables in the p -adic case]. *Matematicheskoe zametki*. 2003;25(1):22–37. Russian. DOI: 10.4213/mzm165.
76. Bernik VI. [A metric theorem on the simultaneous approximation of a zero by the values of integral polynomials]. *Izvestiya Akademii nauk SSSR. Seriya matematicheskaya*. 1980;44(1):24–45. Russian.
77. Zeludevich F. Simultaneous diophantische Approximationen abhängiger Größen in mehreren Metriken. *Acta Arithmetica*. 1986; 46(3):285–296. DOI: 10.4064/aa-46-3-285-296.
78. Bernik V, Budarina N, Dickinson D. A divergent Khintchine theorem in the real, complex, and p -adic fields. *Lithuanian Mathematical Journal*. 2008;48(2):158–173. DOI: 10.1007/s10986-008-9005-9.
79. Budarina N, Dickinson D, Bernik V. Simultaneous Diophantine approximation in the real, complex and p -adic fields. *Mathematical Proceedings of the Cambridge Philosophical Society*. 2010;149(2):193–216. DOI: 10.1017/s0305004110000162.
80. Dombrovskii IR. [Simultaneous approximation of real numbers by algebraic numbers of bounded degree]. *Doklady Akademii nauk BSSR*. 1989;33(3):205–208. Russian.
81. Beresnevich V. Rational points near manifolds and metric Diophantine approximation. *Annals of Mathematics*. 2012;175(1): 187–235. DOI: 10.4007/annals.2012.175.1.5.
82. Beresnevich VV, Velani SL. Simultaneous inhomogeneous Diophantine approximation on manifolds. *Journal of Mathematical Sciences*. 2012;180(5):531–541. DOI: 10.1007/s10958-012-0651-4.
83. Budarina N, Dickinson D, Levesley J. Simultaneous Diophantine approximation on polynomial curves. *Mathematika*. 2010; 56(1):77–85. DOI: 10.1112/s0025579309000382.
84. Budarina N. On a problem of Bernik, Kleinbock and Margulis. *Glasgow Mathematical Journal*. 2011;53(3):669–681. DOI: 10.1017/s0017089511000255.
85. Budarina N, Dickinson D. Simultaneous Diophantine approximation in two metrics and the distance between conjugate algebraic numbers in $\mathbb{R} \times \mathbb{Q}_p$. *Indagationes Mathematicae*. 2012;23(1–2):32–41. DOI: 10.1016/j.indag.2011.09.012.
86. Bernik VI, Borbat VN. [Joint approximation of zero by values of integer-valued polynomials]. *Trudy Matematicheskogo instituta imeni V. A. Steklova*. 1997;218:58–73. Russian.
87. Bernik VI, Budarina N, O’Donnell H. On regular systems of real algebraic numbers of third degree in short intervals. *Sovremennye problemy matematiki*. 2013;17:61–75. DOI: 10.4213/spm43.
88. Bugeaud Y, Mignotte M. Polynômes à coefficients entiers prenant des valeurs positives aux points réels. *Bulletin mathématique de la Société des Sciences Mathématiques de Roumanie*. 2010;53(3):219–224.
89. Bernik V, Götze F. A new connection between metric theory of Diophantine approximations and distribution of algebraic numbers. *Contemporary Mathematics*. 2015;631:33–45. DOI: 10.1090/conm/631/12594.
90. Budarina NV. [Simultaneous Diophantine approximation with non-monotonic right-hand sides]. *Doklady Akademii nauk*. 2011; 437(4):441–443. Russian.
91. Bernik V, Mc Guire S. How small can polynomials be in an interval of given length? *Glasgow Mathematical Journal*. 2020; 62(2):261–280. DOI: 10.1017/S0017089519000077.
92. Bugeaud Y, Mignotte M. Polynomial root separation. *International Journal of Number Theory*. 2010;6(3):587–602. DOI: 10.1142/s1793042110003083.
93. Beresnevich V, Bernik V, Götze F. The distribution of close conjugate algebraic numbers. *Compositio Mathematica*. 2010;146(5): 1165–1179. DOI: 10.1112/S0010437X10004860.
94. Bernik V, Budarina N, O’Donnell H. Discriminants of polynomials in the Archimedean and non-Archimedean metrics. *Acta Mathematica Hungarica*. 2018;154(2):265–278. DOI: <https://doi.org/10.1007/s10474-018-0794-y>.
95. Huxley MN. The rational points close to a curve. *Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Série 4*. 1994;21(3):357–375.
96. Koleda DV. On the asymptotic distribution of algebraic numbers with growing naive height. *Chebyshevskii sbornik*. 2015; 16(1):191–204. Russian. DOI: 10.22405/2226-8383-2015-16-1-191-204.
97. Götze F, Koleda D, Zaporozhets D. Joint distribution of conjugate algebraic numbers: a random polynomial approach. *Advances in Mathematics*. 2020;359:106849. DOI: 10.1016/j.aim.2019.106849.
98. Lebed VV, Bernik VI. Algebraic points on a plane. *Fundamental’naya i prikladnaya matematika*. 2005;11(6):73–80. Russian.



99. Koleda DU. Distribution of the real algebraic numbers of second degree. *Proceedings of the National Academy of Sciences of Belarus. Physics and Mathematics Series*. 2013;3:54–63. Russian.
100. Koleda D. On the density function of the distribution of real algebraic numbers. *Journal de Théorie des Nombres de Bordeaux*. 2017;29(1):179–200. DOI: 10.5802/jtnb.975.
101. Bernik VI, Götze F, Kalosha NI. Counting algebraic numbers in short intervals with rational points. *Journal of the Belarusian State University. Mathematics and Informatics*. 2019;1:4–11. Russian. DOI: 10.33581/2520-6508-2019-1-4-11.
102. Budarina NV, Dickinson D, Bernik VI. Lower bounds for the number of vectors with algebraic coordinates near smooth surfaces. *Doklady of the National Academy of Sciences of Belarus*. 2020;64(1):7–12. Russian. DOI: 10.29235/1561-8323-2020-64-1-7-12.
103. Bernik VI, Götze F. Distribution of real algebraic numbers of arbitrary degree in short intervals. *Izvestiya: Mathematics*. 2015;79(1):18–39. DOI: 10.1070/im2015v079n01abeh002732.
104. Budarina N, Dickinson D. Diophantine approximation on non-degenerate curves with non-monotonic error function. *Bulletin of the London Mathematical Society*. 2009;41(1):137–146. DOI: 10.1112/blms/bdn116.
105. Budarina N. Diophantine approximation on the curves with non-monotonic error function in the p -adic case. *Chebyshevskii sbornik*. 2010;11(1):74–80.
106. Badziahin D, Beresnevich V, Velani S. Inhomogeneous theory of dual Diophantine approximation on manifolds. *Advances in Mathematics*. 2013;232(1):1–35. DOI: 10.1016/j.aim.2012.09.022.
107. Badziahin D. Inhomogeneous Diophantine approximation on curves and Hausdorff dimension. *Advances in Mathematics*. 2010;223(1):329–351. DOI: 10.1016/j.aim.2009.08.005.
108. Bernik V, Dickinson H, Yuan J. Inhomogeneous Diophantine approximation on polynomials in \mathbb{Q}_p . *Acta Arithmetica*. 1999;90(1):37–48. DOI: 10.4064/aa-90-1-37-48.
109. Beresnevich V, Ganguly A, Ghosh A, Velani S. Inhomogeneous dual Diophantine approximation on affine subspaces. *International Mathematics Research Notices*. 2020;12:3582–3613. DOI: 10.1093/imrn/rny124.
110. Beresnevich VV, Vaughan RC, Velani SL. Inhomogeneous Diophantine approximation on planar curves. *Mathematische Annalen*. 2011;349(4):929–942. DOI: 10.1007/s00208-010-0548-9.
111. Arnold VI. [Small denominators and problem of stability of motion in classical and celestial mechanics]. *Uspekhi matematicheskikh nauk*. 1963;18(6):91–192. Russian.
112. Ptashnik BI. *Nekorrektnye granichnye zadachi dlya differentsial'nykh uravnenii s chastnymi proizvodnymi* [Incorrectly defined boundary problems for partial differential equations]. Kyiv: Naukova dumka; 1984. 264 p. Russian.
113. Cadambe VR, Jafar SA. Interference alignment and degrees of freedom of the K -user interference channel. *IEEE Transactions on Information Theory*. 2008;54(8):3425–3441. DOI: 10.1109/TIT.2008.926344.
114. Jafar SA. Interference alignment: a new look at signal dimensions in a communication network. *Foundations and Trends in Communications and Information Theory*. 2011;7(1):1–134. DOI: 10.1561/01000000047.
115. Jafar SA, Shamai Sh. Degrees of freedom region of the MIMO X channel. *IEEE Transactions on Information Theory*. 2008;54(1):151–170. DOI: 10.1109/tit.2007.911262.
116. Beresnevich V, Velani S. Number theory meets wireless communications: an introduction for dummies like us. In: Beresnevich V, Burr A, Nazer B, Velani S, editors. *Number theory meets wireless communications*. [S. l.]: Springer International Publishing; 2020. p. 1–67. DOI: 10.1007/978-3-030-61303-7_1.
117. Budarina N, O'Donnell H. On a problem of Nesterenko: when is the closest root of a polynomial a real number? *International Journal of Number Theory*. 2012;8(3):801–811. DOI: 10.1142/s1793042112500455.
118. Motahari AS, Oveis-Gharan S, Maddah-Ali MA, Khandani AK. Real interference alignment: exploiting the potential of single antenna systems. *IEEE Transactions on Information Theory*. 2014;60(8):4799–4810. DOI: 10.1109/tit.2014.2329865.
119. Sliesoraitienė R. Malerio – Sprindžiuko teoremos analogas trečio laipsnio dviejų kintamųjų polinomams. (II). *Lietuvos matematikos rinkinys*. 1970;10:791–814. Russian.
120. Manstavičius E. Jonas Kubilius 1921–2011. *Lithuanian Mathematical Journal*. 2021;61:285–288. DOI: 10.1007/s10986-021-09522-z.

Received 14.09.2021 / revised 21.10.2021 / accepted 09.11.2021.